

Circular Peaks and Hilbert Series

Pierre Bouchard^a

Jun Ma^{b*}

Yeong-Nan Yeh^{c,†}

June 5, 2008

^a Dépt. de mathématiques, Université du Québec à Montréal, C.P. 8888, Succ. Centre-Ville Montréal,
Canada H3C 3P8

^{b,c} Institute of Mathematics, Academia Sinica, Taipei, Taiwan

Abstract

The circular peak set of a permutation σ is the set $\{\sigma(i) \mid \sigma(i-1) < \sigma(i) > \sigma(i+1)\}$. Let $\mathcal{P}_n$ be the set of all the subset $S \subseteq [n]$ such that there exists a permutation σ which has the circular set S . We can make the set $\mathcal{P}_n$ into a poset $\mathcal{P}_n$ by defining $S \preceq T$ if $S \subseteq T$ as sets. In this paper, we prove that the poset $\mathcal{P}_n$ is a simplicial complex on the vertex set $[3, n]$. We study the f -vector, the f -polynomial, the reduced Euler characteristic, the Möbius function, the h -vector and the h -polynomial of $\mathcal{P}_n$. We also derive the zeta polynomial of $\mathcal{P}_n$ and give the formula for the number of the chains in $\mathcal{P}_n$. By the poset $\mathcal{P}_n$, we define two algebras $\mathcal{A}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n}$. We consider the Hilbert polynomials and the Hilbert series of the algebra $\mathcal{A}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n}$.

Keyword: Circular peak; Hilbert Polynomial; Hilbert series; Permutation; Poset; Simplicial complex;

*Email address of the corresponding author: majun@math.sinica.edu.tw

†Partially supported by NSC 96-2115-M-006-012

1 Introduction

Throughout this paper, let $[m, n] := \{m, m+1, \dots, n\}$, $[n] := [1, n]$ and $\mathfrak{S}_n$ be the set of all the permutations in the set $[n]$. We will write permutations of $\mathfrak{S}_n$ in the form $\sigma = (\sigma(1)\sigma(2)\cdots\sigma(n))$.

We say that a permutation σ has a circular descent of value $\sigma(i)$ if $\sigma(i) > \sigma(i+1)$ for any $i \in [n-1]$. The circular descent set of a permutation σ , denoted $CDES(\sigma)$, is the set $\{\sigma(i) \mid \sigma(i) > \sigma(i+1)\}$. For any $S \subseteq [n]$, define a set $CDES_n(S)$ as $CDES_n(S) = \{\sigma \in \mathfrak{S}_n \mid CDES(\sigma) = S\}$ and use $cdes_n(S)$ to denote the number of the permutations in the set $CDES_n(S)$, i.e., $cdes_n(S) = |CDES_n(S)|$. In a join work [1], Hungyung Zhang et al. derive the explicit formula for $cdes_n(S)$. As a application of the main results in [1], they also give the enumeration of permutation tableaux according to their shape and generalizes the results in [3]. Moreover, Robert J. Clarke et al. [2] gave the conceptions of linear peak and cyclic peak and studied some new Mahonian permutation statistics. In this paper, we are interested in the circular peaks of permutations. A permutation σ has a *circular peak* of value $\sigma(i)$ if $\sigma(i-1) < \sigma(i) > \sigma(i+1)$ for any $i \in [2, n-1]$. The *circular peak set* of a permutation σ , denoted $CP(\sigma)$, is the set $\{\sigma(i) \mid \sigma(i-1) < \sigma(i) > \sigma(i+1)\}$. For example, the circular peak set of $\sigma = (48362517)$ is $\{5, 6, 8\}$. Since σ has no circular peaks when $n \leq 2$, we may always suppose that $n \geq 3$. For any $S \subseteq [n]$, define a set $CP_n(S)$ as $CP_n(S) = \{\sigma \in \mathfrak{S}_n \mid CP(\sigma) = S\}$. Obviously, if $\{1, 2\} \subseteq S$, then $CP_n(S) = \emptyset$.

Example 1.1

$$CP_5(\{4, 5\}) = \{ \quad 14253, 14352, 24153, 34152, 24351, 34251, \\ 15243, 15342, 25143, 35142, 25341, 35241 \quad \}$$

Suppose that $S = \{i_1, i_2, \dots, i_k\}$, where $i_1 < i_2 < \dots < i_k$. We prove that the necessary and sufficient conditions for $CP_n(S) \neq \emptyset$ are $i_j \geq 2j+1$ for all $j \in [k]$.

Let $\mathcal{P}_n = \{S \mid CP_n(S) \neq \emptyset\}$. we can make the set $\mathcal{P}_n$ into a poset $\mathcal{P}_n$ by defining $S \preceq T$ if $S \subseteq T$ as sets. We draw the Hasse diagrams of $\mathcal{P}_3$, $\mathcal{P}_4$ and $\mathcal{P}_5$ as follows.




Fig.1. the Hasse diagrams of $\mathcal{P}_3$, $\mathcal{P}_4$ and $\mathcal{P}_5$

There is a very interesting results: $\mathcal{P}_n$ is a simplicial complex on the vertex set $[3, n]$. It is easy to obtain that the dimension of the simplicial complex $\mathcal{P}_n$ is $\lfloor \frac{1}{2}(n-1) \rfloor - 1$. As we will see, the number of the elements in $\mathcal{P}_n$ involves in the left factors of Dyck paths of length $n-1$, counted by the $(n-1)$ -th central binomial coefficients b_{n-1} (see Cori and Viennot [4]), where $b_{n-1} = \binom{n-1}{\lfloor \frac{1}{2}(n-1) \rfloor}$; the number $p_{n,i}$ of the faces of dimension i in $\mathcal{P}_n$ equals the number of the left factors of Dyck paths from $(0,0)$ to $(n-1, n-2i-3)$, counted by $b_{n-1,i+1} = \frac{n-2i-2}{i+1} \binom{n-1}{i}$ [4].

We derive the recurrence relations for the poset $\mathcal{P}_n$: $\mathcal{P}_{n+1} \cong \mathbf{2} \times \mathcal{P}_n$ if n is even; $\mathcal{P}_{n+1} \cong (\mathbf{2} \times \mathcal{P}_n) \setminus (\{1\} \times \mathcal{P}_{n, \lfloor \frac{1}{2}(n-1) \rfloor - 1})$ if n is odd, where the notation $\mathbf{n}$ denotes a poset formed by the set $[n]$ with its usual order.

It is very important to obtain the f -vector, the f -polynomial and the reduced Euler characteristic of a simplicial complex. The integral sequence $(p_{n,-1}, p_{n,0}, \dots, p_{n, \lfloor \frac{1}{2}(n-1) \rfloor - 1})$ is called the f -vector of $\mathcal{P}_n$. The f -polynomial of $\mathcal{P}_n$ is defined to be the polynomial $\mathcal{P}_n(x) = \sum_{i=0}^{\lfloor \frac{1}{2}(n-1) \rfloor} p_{n,i-1} x^{\lfloor \frac{1}{2}(n-1) \rfloor - i}$. We give the recurrence relations for the f -vector and the f -polynomial of $\mathcal{P}_n$. Let $\mathcal{P}(x, y) = \sum_{n \geq 3} \mathcal{P}_n(x) y^n$, then $\mathcal{P}(x, y) = \frac{xy^2(x+2) - xy^2 C(y^2)}{x - (x+1)y^2} \frac{1+y+xy}{x+1} - y^2$, where $C(y) = \frac{1-\sqrt{1-4y}}{2y}$. The reduced Euler characteristic $\tilde{\chi}(\mathcal{P}_n)$ of $\mathcal{P}_n$ satisfies that $\tilde{\chi}(\mathcal{P}_n) = 0$ if n is odd; $\tilde{\chi}(\mathcal{P}_n) = \frac{2(-1)^{\frac{n}{2}}}{n} \binom{n-2}{\frac{1}{2}(n-2)}$ if n is even. As a corollary, the zeta poly-

nomial of $\mathcal{P}_n$ is $(m-1)^{\lfloor \frac{n-1}{2} \rfloor} \mathcal{P}_n(\frac{1}{m-1})$, which is the number of multichains $S_{n,1} \preceq S_{n,2} \preceq \cdots \preceq S_{n,m-1}$ in $\mathcal{P}_n$ for any $m \geq 2$. The number of the chains $S_{n,1} \prec S_{n,2} \prec \cdots \prec S_{n,i}$ in $\mathcal{P}_n$ equals

$$\sum \binom{n}{d_1, d_2, \dots, d_{i+1}} \frac{2d_{i+1} - n}{n},$$

where the sum is over all $(d_1, \dots, d_{i+1})$ such that $\sum_{k=1}^{i+1} d_k = n$, $d_1 \geq 0$, $d_k \geq 1$ for all $2 \leq k \leq i$ and $d_{i+1} \geq n - \lfloor \frac{n-1}{2} \rfloor$.

We are interested in the h -vector and the h -polynomial of the simplicial complex $\mathcal{P}_n$. We obtain the explicit formula for h -vector and give the recurrence relations for the h -vector and the h -polynomial of $\mathcal{P}_n$.

We fix a field $\mathbb{K}$ and let $m = \binom{n-1}{\lfloor \frac{1}{2}(n-1) \rfloor}$ and all the elements of $\mathcal{P}_n$ be $S_{n,1}, S_{n,2}, \dots, S_{n,m}$ for any $n \geq 3$. Let $\mathcal{I}_{\mathcal{P}_n}$ be the ideal of the polynomial ring $\mathbb{K}[x_1, \dots, x_m]$ which is generated by all polynomials $x_i x_j$ such that $S_{n,i}$ and $S_{n,j}$ are incomparable in the poset $\mathcal{P}_n$. Let $\mathcal{J}_{\mathcal{P}_n}$ be the ideal of the polynomial ring $\mathbb{K}[x_1, \dots, x_m]$ which is generated by all polynomials $x_i x_j$ such that $S_{n,i}$ and $S_{n,j}$ are incomparable in $\mathcal{P}_n$ and the polynomials x_i^2 for all $i \in [m]$. Define the two algebras $\mathcal{A}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n}$ as the quotients $\mathcal{A}_{\mathcal{P}_n} = \mathbb{K}[x_1, \dots, x_m] / \mathcal{I}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n} = \mathbb{K}[x_1, \dots, x_m] / \mathcal{J}_{\mathcal{P}_n}$, respectively. We study the Hilbert polynomials and the Hilbert series of the algebras $\mathcal{A}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n}$.

This paper is organized as follows. In Section 2, we give the necessary and sufficient conditions for $CP_n(S) \neq \emptyset$. In Section 3, we prove that the poset $\mathcal{P}_n$ is a simplicial complex and study its properties. In Section 4, we define the two algebras $\mathcal{A}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n}$ and consider their Hilbert polynomial and Hilbert series.

2 The Necessary and Sufficient Conditions for $CP_n(S) \neq \emptyset$

$\emptyset$

In this section, we will give the necessary and sufficient conditions for $CP_n(S) \neq \emptyset$ for any $n \geq 3$ and $S \subseteq [n]$.

Theorem 2.1 *Suppose that n is an integer with $n \geq 3$. Let $S = \{i_1, i_2, \dots, i_k\}$ be a subset of $[n]$, where $i_1 < i_2 < \dots < i_k$. Then the necessary and sufficient conditions for $CP_n(S) \neq \emptyset$ are $i_j \geq 2j + 1$ for all $j \in [k]$.*

Proof. First, we suppose that $CP_n(S) \neq \emptyset$ and let $\sigma \in CP_n(S)$. For any $j \in [k]$, since all the numbers $i_1, i_2, \dots, i_j$ are peaks of σ , we have $i_j - j \geq j + 1$, hence, $i_j \geq 2j + 1$.

Conversely, when $i_j \geq 2j + 1$ for all $j \in [k]$, let $T = [i_k] \setminus S$, then the set T has at least $k + 1$ elements. So, suppose that $T = \{a_1, a_2, \dots, a_m\}$ with $a_1 < a_2 < \dots < a_m$. We consider $\sigma = a_1 i_1 a_2 i_2 \dots a_k i_k a_{k+1} \dots a_m (i_k + 1) \dots n$. Obviously, $CP(\sigma) \subseteq S$. If $CP(\sigma) \neq S$, then there exists a minimal $j \in [k]$ such that i_j is not a circular peak of σ . So, $a_{j+1} > i_j$. This implies that $i_j = 2j$, a contradiction. Hence, $CP(\sigma) = S$ and $CP_n(S) \neq \emptyset$. ■

Corollary 2.1 *Suppose that n is an integer with $n \geq 3$ and $S \subseteq [n]$. If $CP_n(S) \neq \emptyset$, then $|S| \leq \lfloor \frac{n-1}{2} \rfloor$.*

Proof. Suppose that $S = \{i_1, i_2, \dots, i_k\}$ with $i_1 < i_2 < \dots < i_k$. Since $CP_n(S) \neq \emptyset$, Theorem 2.1 tells us that $i_k \geq 2k + 1$. $i_k \leq n$ implies that $k \leq \lfloor \frac{n-1}{2} \rfloor$. ■

Corollary 2.2 *Let $n \geq 3$ and $S \subseteq [n]$. Suppose that $CP_n(S) \neq \emptyset$. Then when $|S| < \lfloor \frac{n-1}{2} \rfloor$, we have $CP_{n+1}(S \cup \{n+1\}) \neq \emptyset$; when $|S| = \lfloor \frac{n-1}{2} \rfloor$, we have $CP_{n+1}(S \cup \{n+1\}) \neq \emptyset$ if n is even; otherwise, $CP_{n+1}(S \cup \{n+1\}) = \emptyset$.*

Proof. Suppose that $k = |S|$. $k < \lfloor \frac{n-1}{2} \rfloor$ implies that $2(k+1) + 1 \leq 2\lfloor \frac{n-1}{2} \rfloor + 1 < n+1$.

So, $CP_{n+1}(S \cup \{n+1\}) \neq \emptyset$ when $|S| < \lfloor \frac{n-1}{2} \rfloor$. For the case $k = \lfloor \frac{n-1}{2} \rfloor$, we have

$$2(k+1) + 1 = \begin{cases} n+1 & \text{if } n \text{ is even} \\ n+2 & \text{if } n \text{ is odd} \end{cases}.$$

By Theorem 2.1, $CP_{n+1}(S \cup \{n+1\}) \neq \emptyset$ if n is even; otherwise, $CP_{n+1}(S \cup \{n+1\}) = \emptyset$. ■

3 The Simplicial Complex $\mathcal{P}_n$

In this section, we will prove that the poset $\mathcal{P}_n$ is a simplicial complex on the vertex set $[3, n]$ for any $n \geq 3$, and then study the properties of the Simplicial Complex $\mathcal{P}_n$. Following [5], we define a *simplicial complex* Δ on a vertex set V as a collection of subsets of V satisfying:

- (1) If $x \in V$, then $\{x\} \in \Delta$, and
- (2) if $S \in \Delta$ and $T \subseteq S$, then $T \in \Delta$.

Theorem 3.1 *Let $n \geq 3$. Then $\mathcal{P}_n$ is a simplicial complex of the set $[3, n]$ and has the dimension $\lfloor \frac{n-1}{2} \rfloor - 1$.*

Proof. Obviously, $\emptyset \in \mathcal{P}_n$. For any $3 \leq x \leq n$, Theorem 2.1 implies that $\{x\} \in \mathcal{P}_n$. Noting that if $CP_n(T) = \emptyset$ then $CP_n(S) = \emptyset$ for any $S \supseteq T$, we have if $S \in \mathcal{P}_n$ and $T \subseteq S$ then $T \in \mathcal{P}_n$. Hence, $\mathcal{P}_n$ is a simplicial complex of the set $[3, n]$. ■

An element $S \in \mathcal{P}_n$ is called a *face* of $\mathcal{P}_n$, and the *dimension* of S is defined to be $|S| - 1$. In particular, the void set $\emptyset$ is always a face of $\mathcal{P}_n$ of dimension -1 . Also define the dimension of $\mathcal{P}_n$ by $\dim \mathcal{P}_n = \max_{S \in \mathcal{P}_n} (\dim S)$.

Theorem 3.2 *The simplicial complex $\mathcal{P}_n$ has the dimension $\lfloor \frac{n-1}{2} \rfloor - 1$.*

Proof. Taking $S = \{3, 5, \dots, 2\lfloor \frac{n-1}{2} \rfloor + 1\}$, by Theorem 2.1, we have $S \in \mathcal{P}_n$. From Corollary 2.1 it follows that the dimension of $\mathcal{P}_n$ is $\lfloor \frac{n-1}{2} \rfloor - 1$. ■

There are very close relations between the number of the elements of $\mathcal{P}_n$ and the number of left factor of Dyck path of length n . An n -Dyck path is a lattice path in the first quadrant starting at $(0, 0)$ and ending at $(2n, 0)$ with only two kinds of steps—*rise step*: $U = (1, 1)$ and *fall step*: $D = (1, -1)$. We can also consider an n -Dyck path P as a word of $2n$ letters using only U and D . Let $L = w_1w_2 \cdots w_n$ be a word, where $w_j \in \{U, D\}$ and $n \geq 0$. If there is another word R which consists of U and D such that LR forms a Dyck path, then L is called an n -left factor of Dyck path. Let $\mathcal{L}_n$ denote the set of all n -left factor of Dyck paths. It is well known that $|\mathcal{L}_n|$, the cardinality of $\mathcal{L}_n$, equals the n th Central binomial number $b_n = \binom{n}{\lfloor \frac{n}{2} \rfloor}$. In the following lemma, we give a bijection ϕ from the sets $\mathcal{P}_n$ to $\mathcal{L}_{n-1}$.

Lemma 3.1 *There is a bijection ϕ between the sets $\mathcal{P}_n$ and $\mathcal{L}_{n-1}$ for any $n \geq 3$. Furthermore, the number of the elements in $\mathcal{P}_n$ is $\binom{n-1}{\lfloor \frac{n-1}{2} \rfloor}$.*

Proof. For any $S \in \mathcal{P}_n$, we construct a word $\phi(S) = w_1w_2 \cdots w_{n-1}$ as follows:

$$w_i = \begin{cases} D & \text{if } i+1 \in S \\ U & \text{if } i+1 \notin S \end{cases}$$

for any $i \in [n-1]$. Theorem 2.1 implies that $\phi(S)$ is a $(n-1)$ -left factor of Dyck path. Conversely, for any a n -left factor of Dyck path $w_1w_2 \cdots w_{n-1}$, let $S = \{i+1 \mid w_i = D\}$, then $CP_n(S) \neq \emptyset$. Hence, the mapping ϕ is a bijection. Note that the number of $(n-1)$ -left factor of Dyck path is $\binom{n-1}{\lfloor \frac{n-1}{2} \rfloor}$. Hence, $|\mathcal{P}_n| = \binom{n-1}{\lfloor \frac{n-1}{2} \rfloor}$. ■

Now, we are in a position to obtain the number $p_{n,i}$ of the faces of dimension i in $\mathcal{P}_n$. For any $i \geq 0$, let $\mathcal{L}_{n,i}$ denote the set of all n -left factor of Dyck paths from $(0, 0)$ to $(n, n-2i)$. Define a set $\mathcal{P}_{n,i}$ as the set of all the faces of dimension i in $\mathcal{P}_n$, i.e., $\mathcal{P}_{n,i} = \{S \in \mathcal{P}_n \mid |S| = i+1\}$ for any $-1 \leq i \leq \lfloor \frac{n-1}{2} \rfloor - 1$. Clearly, $p_{n,i} = |\mathcal{P}_{n,i}|$.

Corollary 3.1 *Let $n \geq 3$. There is a bijection between the sets $\mathcal{P}_{n,i}$ and $\mathcal{L}_{n-1,i+1}$ for any*

$-1 \leq i \leq \lfloor \frac{n-1}{2} \rfloor - 1$. Furthermore, we have

$$p_{n,i} = \begin{cases} 1 & \text{if } i = -1 \\ \frac{n-2i-2}{i+1} \binom{n-1}{i} & \text{if } 0 \leq i \leq \lfloor \frac{n-1}{2} \rfloor - 1 \end{cases}$$

Proof. We just consider the case with $i \geq 0$. For any $S \in \mathcal{P}_{n,i}$, $|S| = i + 1$ implies that the number of letter D in the word $\phi(S)$ is $i + 1$. Hence, $\phi(S)$ is a left factor of Dyck path from $(0, 0)$ to $(n - 1, n - 2i - 3)$. So, $\phi(S) \in \mathcal{L}_{n-1,i+1}$. ■

If P and Q are posets, then the *direct product* of P and Q is the poset $P \times Q$ on the set $\{(x, y) \mid x \in P \text{ and } y \in Q\}$ such that $(x, y) \leq (x', y')$ in $P \times Q$ if $x \leq x'$ in P and $y \leq y'$ in Q . Recall that the poset $\mathbf{n}$ is formed by the set $[n]$ with its usual order. By Corollary 2.2, we obtain a method for constructing the poset $\mathcal{P}_{n+1}$ from $\mathcal{P}_n$.

Theorem 3.3 $\mathcal{P}_{n+1} \cong \mathbf{2} \times \mathcal{P}_n$ if n is even; $\mathcal{P}_{n+1} \cong (\mathbf{2} \times \mathcal{P}_n) \setminus (\{1\} \times \mathcal{P}_{n, \lfloor \frac{n-1}{2} \rfloor - 1})$ if n is odd.

By Theorem 3.3, it is easy for us to obtain the Möbius function, the recurrence relations for the f -vector and the f -polynomial of the poset $\mathcal{P}_n$.

Corollary 3.2 Let $\mu_{\mathcal{P}_n}$ be the Möbius function of the poset $\mathcal{P}_n$. Then

$$\mu_{\mathcal{P}_n}(S, T) = (-1)^{|T| - |S|}$$

for any $S \preceq T$ in $\mathcal{P}_n$.

Proof. Obviously, $\mu_{\mathcal{P}_3}(\emptyset, \{3\}) = -1$. By induction for n , we assume that $\mu_{\mathcal{P}_n}(S, T) = (-1)^{|T| - |S|}$ for any $S \preceq T$ in $\mathcal{P}_n$. Lemma 3.3 tells us that $\mathcal{P}_{n+1} \cong \mathbf{2} \times \mathcal{P}_n$ if n is even; $\mathcal{P}_{n+1} \cong (\mathbf{2} \times \mathcal{P}_n) \setminus (\{1\} \times \mathcal{P}_{n, \lfloor \frac{n-1}{2} \rfloor})$ if n is odd. We conclude from the product theorem that

$$\mu_{\mathcal{P}_{n+1}}(S, T) = \begin{cases} \mu_{\mathcal{P}_n}(S \setminus \{n+1\}, T \setminus \{n+1\}) & \text{if } n+1 \in S \cap T \\ \mu_{\mathcal{P}_n}(S, T) & \text{if } n+1 \notin S \cup T \\ -\mu_{\mathcal{P}_n}(S, T \setminus \{n+1\}) & \text{if } n+1 \notin S \text{ and } n+1 \in T \end{cases}$$

for any $S \prec T$. Note that $\mu_{\mathcal{P}_n}(S \setminus \{n+1\}, T \setminus \{n+1\}) = (-1)^{|T|-1-(|S|-1)} = (-1)^{|T|-|S|}$ and $-\mu_{\mathcal{P}_n}(S, T \setminus \{n+1\}) = -(-1)^{|T|-1-|S|} = (-1)^{|T|-|S|}$. Hence, $\mu_{\mathcal{P}_{n+1}}(S, T) = (-1)^{|T|-|S|}$. ■

Corollary 3.3 *Let $n \geq 3$. The sequence $p_{n,i}$ satisfies the following recurrence relation: when n is even ,*

$$p_{n+1,i} = \begin{cases} p_{n,i} & \text{if } i = -1 \\ p_{n,i-1} + p_{n,i} & \text{if } i = 0, 1, \dots, \frac{n}{2} - 2 \\ p_{n,i-1} & \text{if } i = \frac{n}{2} - 1 \end{cases} ;$$

when n is odd ,

$$p_{n+1,i} = \begin{cases} p_{n,i} & \text{if } i = -1 \\ p_{n,i-1} + p_{n,i} & \text{if } i = 0, 1, \dots, \frac{n-3}{2} \end{cases}$$

with initial conditions $(p_{3,-1}, p_{3,0}) = (1, 1)$.

Proof. First, we consider the case with n even. It is easy to check that $p_{n+1,-1} = p_{n,-1} = 1$. For any $S \in \mathcal{P}_{n+1, \frac{1}{2}n-1}$, Corollary 2.2 tells us that $n+1 \in S$. Note that $S \in \mathcal{P}_{n+1, \frac{1}{2}n-1}$ if and only if $S \setminus \{n+1\} \in \mathcal{P}_{n, \frac{1}{2}n-2}$. Hence, $p_{n+1, \frac{1}{2}n-1} = p_{n, \frac{1}{2}n-2}$. Let $0 \leq i \leq \frac{1}{2}n - 2$, obviously, $\mathcal{P}_{n,i} \subseteq \mathcal{P}_{n+1,i}$. For any $S \in \mathcal{P}_{n+1,i}$ with $n+1 \in S$, $S \setminus \{n+1\}$ can be viewed as a element of $\mathcal{P}_{n,i-1}$. Conversely, for any $S \in \mathcal{P}_{n,i-1}$, Corollary 2.2 implies that $S \cup \{n+1\} \in \mathcal{P}_{n+1,i}$. Hence, $p_{n+1,i} = p_{n,i-1} + p_{n,i}$. Similarly, we can consider the case with n odd. ■

Theorem 3.4 *The f -polynomial $\mathcal{P}_n(x)$ of the simplicial complex $\mathcal{P}_n$ satisfy the following recurrence relation: when n is even, then*

$$\mathcal{P}_{n+1}(x) = (1+x)\mathcal{P}_n(x),$$

and when n is odd, then

$$x\mathcal{P}_{n+1}(x) = (1+x)\mathcal{P}_n(x) - \frac{2}{n+1} \binom{n-1}{\frac{n-1}{2}}.$$

for any $n \geq 3$, with initial condition $\mathcal{P}_3(x) = x + 1$.

Let $\mathcal{P}(x, y) = \sum_{n \geq 3} \mathcal{P}_n(x) y^n$. Then

$$\mathcal{P}(x, y) = \frac{xy^2(x+2) - xy^2C(y^2)}{x - (x+1)y^2} \frac{1+y+xy}{x+1} - y^2,$$

where $C(y) = \frac{1-\sqrt{1-4y}}{2y}$.

Proof. Obviously, $\mathcal{P}_3(x) = x+1$. When n is odd, we suppose that $n = 2i+1$ with $i \geq 1$.

Corollary 3.3 implies that

$$x\mathcal{P}_{2i+2}(x) = (1+x)\mathcal{P}_{2i+1}(x) - \frac{1}{(i+1)} \binom{2i}{i}.$$

Similarly, when n is even, we suppose that $n = 2i$ with $i \geq 2$. By corollary 3.3, we have

$$\mathcal{P}_{2i+1}(x) = (1+x)\mathcal{P}_{2i}(x).$$

Let $\mathcal{P}_{odd}(x, y) = \sum_{i \geq 1} \mathcal{P}_{2i+1}(x) y^{2i+1}$ and $\mathcal{P}_{even}(x, y) = \sum_{m \geq 2} \mathcal{P}_{2i}(x) y^{2i}$, then $\mathcal{P}_{odd}(x, y) = (x+1)y^3 + (x+1)y\mathcal{P}_{even}(x, y)$ and $\mathcal{P}(x, y) = \mathcal{P}_{odd}(x, y) + \mathcal{P}_{even}(x, y)$. It is easy to check that $x\mathcal{P}_{2i+3}(x) = (1+x)^2\mathcal{P}_{2i+1}(x) - \frac{1}{i+1} \binom{2i}{i}x$. So, $\mathcal{P}_{odd}$ satisfies the following equation

$$x\mathcal{P}_{odd}(x, y) = (x+1)^2y^2\mathcal{P}_{odd}(x, y) + x(x+2)y^3 - xy^3C(y^2),$$

where $C(y) = \frac{1-\sqrt{1-4y}}{2y}$. Equivalently,

$$\mathcal{P}_{odd}(x, y) = \frac{xy^3(x+2) - xy^3C(y^2)}{x - (x+1)y^2}.$$

Hence,

$$\mathcal{P}(x, y) = \frac{xy^2(x+2) - xy^2C(y^2)}{x - (x+1)y^2} \frac{1+y+xy}{x+1} - y^2.$$

■

Define the *reduced Euler characteristic* of $\mathcal{P}_n$ by $\tilde{\chi}(\mathcal{P}_n) = \sum_{i=0}^{\lfloor \frac{1}{2}(n-1) \rfloor} (-1)^{i-1} p_{n,i-1}$.

Corollary 3.4 For any $n \geq 3$, $\tilde{\chi}(\mathcal{P}_n) = \begin{cases} 0 & \text{if } n \text{ is odd} \\ \frac{2(-1)^{\frac{n}{2}}}{n} \binom{n-2}{\frac{1}{2}(n-2)} & \text{if } n \text{ is even} \end{cases}$

Proof. Clearly, $\mathcal{P}_3(-1) = 0$. Theorem 3.4 tells us that

$$\mathcal{P}_{n+1}(-1) = \begin{cases} 0 & \text{if } n \text{ is even} \\ \frac{2}{n+1} \binom{n-1}{\frac{1}{2}(n-1)} & \text{if } n \text{ is odd} \end{cases}$$

for any $n \geq 4$. Since $\tilde{\chi}(\mathcal{P}_n) = (-1)^{\lfloor \frac{n-1}{2} \rfloor - 1} \mathcal{P}_n(-1)$, we have

$$\tilde{\chi}(\mathcal{P}_n) = \begin{cases} 0 & \text{if } n \text{ is odd} \\ \frac{2(-1)^{\frac{n}{2}-2}}{n} \binom{n-2}{\frac{1}{2}(n-2)} & \text{if } n \text{ is even} \end{cases}$$

■

Let P be a finite post. Define $Z(P, i)$ to be the number of multichains $x_1 \leq x_2 \leq \cdots \leq x_{i-1}$ in P for any $i \geq 2$. $Z(P, i)$ is called the *zeta polynomial* of P . We state Proposition 3.11.1a and Proposition 3.14.2 in [5] as the following lemma.

Lemma 3.2 *Suppose that P is a poset. (1) Let d_i be the number of chains $x_1 < x_2 < \cdots < x_{i-1}$ in P . Then $Z(P, i) = \sum_{j \geq 2} d_j \binom{i-2}{j-2}$.*

(2) If P is simplicial and graded, then $Z(P, x+1)$ is the rank-generating function of P .

Corollary 3.5 *Let $n \geq 3$ and $m \geq 2$. Then $Z(\mathcal{P}_n, i) = (i-1)^{\lfloor \frac{n-1}{2} \rfloor} \mathcal{P}_n(\frac{1}{i-1})$ for any $i \geq 2$. Furthermore, $Z(\mathcal{P}_n, i)$ satisfies the recurrence relations: $Z(\mathcal{P}_{n+1}, i) = iZ(\mathcal{P}_n, i) - \varepsilon(n) \frac{2(i-1)^{\frac{1}{2}(n+1)}}{n+1} \binom{n-1}{\frac{1}{2}(n-1)}$, where $\varepsilon(n) = 0$ if n is even; $\varepsilon(n) = 1$ otherwise, with initial condition $Z(\mathcal{P}_3, i) = i$.*

Proof. Let $\mathcal{P}_n(x)$ be the f -polynomial of $\mathcal{P}_n$, then the rank-generating function of $\mathcal{P}_n$ is $x^{\lfloor \frac{1}{2}(n-1) \rfloor} \mathcal{P}_n(\frac{1}{x})$. Lemma 3.2(2) implies that $Z(\mathcal{P}_n, i) = (i-1)^{\lfloor \frac{n-1}{2} \rfloor} \mathcal{P}_n(\frac{1}{i-1})$. The recurrence relations for $Z(\mathcal{P}_n, i)$ follows Theorem 3.4. ■

Let $d_{\mathcal{P}_n, i}$ be the number of the chains $S_{n,1} \prec S_{n,2} \prec \cdots \prec S_{n,i}$ of $\mathcal{P}_n$.

Theorem 3.5 *For any $i \geq 1$,*

$$d_{\mathcal{P}_n, i} = \sum \binom{n}{d_1, d_2, \dots, d_{i+1}} \frac{2d_{i+1} - n}{n},$$

where the sum is over all $(d_1, \dots, d_{i+1})$ such that $\sum_{k=1}^{i+1} d_k = n$, $d_1 \geq 0$, $d_k \geq 1$ for all $2 \leq k \leq i$ and $d_{i+1} \geq n - \lfloor \frac{n-1}{2} \rfloor$.

Proof. Let $i \geq 1$ and $S_{n,1} \prec S_{n,2} \prec \dots \prec S_{n,i}$ be a chain of $\mathcal{P}_n$. Suppose that $|S_{n,k}| = j_k$ for any $k \in [i]$, then $0 \leq j_1 < j_2 < \dots < j_i \leq \lfloor \frac{n-1}{2} \rfloor$. There are p_{n,j_i-1} ways to obtain the set $S_{n,i}$. Given $S_{n,k}$ with $k \geq 2$, there are $\binom{j_k}{j_{k-1}}$ ways to form the subset $S_{n,k-1} \subseteq S_{n,k}$. Hence,

$$\begin{aligned} d_{\mathcal{P}_n,i} &= \sum_{0=j_0 \leq j_1 < j_2 < \dots < j_i \leq \lfloor \frac{n-1}{2} \rfloor} \prod_{k=0}^{i-1} \binom{j_{k+1}}{j_k} p_{n,j_i-1} \\ &= \sum \binom{n}{d_1, d_2, \dots, d_{i+1}} \frac{2d_{i+1} - n}{n}, \end{aligned}$$

where the sum is over all $(d_1, \dots, d_{i+1})$ such that $\sum_{k=1}^{i+1} d_k = n$, $d_1 \geq 0$, $d_k \geq 1$ for all $2 \leq k \leq i$ and $d_{i+1} \geq n - \lfloor \frac{n-1}{2} \rfloor$. ■

Corollary 3.6 For any $n \geq 3$,

$$\mathcal{P}_n(x) = \sum_{i=2}^{\lfloor \frac{n-1}{2} \rfloor + 2} \frac{x^{\lfloor \frac{n-1}{2} \rfloor + 2 - i}}{(i-2)!} \prod_{j=1}^{i-2} (1 - jx) \sum \binom{n}{d_1, d_2, \dots, d_i} \frac{2d_i - n}{n}$$

where the second sum is over all $(d_1, \dots, d_i)$ such that $\sum_{k=1}^i d_k = n$, $d_1 \geq 0$, $d_k \geq 1$ for all $2 \leq k \leq i-1$ and $d_i \geq n - \lfloor \frac{n-1}{2} \rfloor$.

Proof. Lemma 3.2(1) implies that $Z(\mathcal{P}_n, i) = \sum_{j=2}^{\lfloor \frac{n-1}{2} \rfloor + 2} d_{\mathcal{P}_n, j-1} \binom{i-2}{j-2}$. By Corollary 3.5, we have

$$\mathcal{P}_n \left(\frac{1}{i-1} \right) = \left(\frac{1}{i-1} \right)^{\lfloor \frac{n-1}{2} \rfloor} \sum_{j=2}^{\lfloor \frac{n-1}{2} \rfloor + 2} d_{\mathcal{P}_n, j-1} \binom{i-2}{j-2}$$

for any $i \geq 2$. Note that

$$x^{\lfloor \frac{n-1}{2} \rfloor} \sum_{j=2}^{\lfloor \frac{n-1}{2} \rfloor + 2} d_{\mathcal{P}_n, j-1} \binom{\frac{1}{x}}{j-2} = \sum_{j=2}^{\lfloor \frac{n-1}{2} \rfloor + 2} \frac{x^{\lfloor \frac{n-1}{2} \rfloor + 2 - j}}{(j-2)!} \prod_{k=1}^{j-2} (1 - kx) d_{\mathcal{P}_n, j-1}$$

is a polynomial. Hence, $\mathcal{P}_n(x) = \sum_{j=2}^{\lfloor \frac{n-1}{2} \rfloor + 2} \frac{x^{\lfloor \frac{n-1}{2} \rfloor + 2 - j}}{(j-2)!} \prod_{k=1}^{j-2} (1 - kx) d_{\mathcal{P}_n, j-1}$. ■

Let $\mathcal{H}_n(x) = \mathcal{P}_n(x-1) = \sum_{i=0}^{\lfloor \frac{1}{2}(n-1) \rfloor} h_{n,i} x^{\lfloor \frac{1}{2}(n-1) \rfloor - i}$, then $\mathcal{H}_n(x)$ is called *h-polynomial* of $\mathcal{P}_n$ and the sequence $(h_{n,0}, h_{n,1}, \dots, h_{n, \lfloor \frac{1}{2}(n-1) \rfloor})$ *h-vector* of $\mathcal{P}_n$.

Corollary 3.7 *The h-polynomial $\mathcal{H}_n(x)$ of the simplicial complex $\mathcal{P}_n$ satisfies the recurrence relation: when n is even,*

$$\mathcal{H}_{n+1}(x) = x\mathcal{H}_n(x),$$

and when n is odd,

$$(x-1)\mathcal{H}_{n+1}(x) = x\mathcal{H}_n(x) - \frac{2}{(n+1)} \binom{n-1}{\frac{n-1}{2}},$$

for any $n \geq 3$, with initial condition $\mathcal{H}_3(x) = x$.

Let $\mathcal{H}(x, y) = \sum_{n \geq 3} \mathcal{H}_n(x) y^n$, then

$$\mathcal{H}(x, y) = \frac{[(x^2 - 1)y^2 - (x-1)y^2 C(y^2)](1+xy)}{x(x-1-xy^2)} - y^2.$$

Furthermore, let $(h_{n,0}, h_{n,1}, \dots, h_{n, \lfloor \frac{n-1}{2} \rfloor})$ be the *h-vector* of $\mathcal{P}_n$, then

$$h_{n,i} = \frac{\lfloor \frac{n}{2} \rfloor - i}{\lfloor \frac{n}{2} \rfloor + i} \binom{\lfloor \frac{n}{2} \rfloor + i}{\lfloor \frac{n}{2} \rfloor}.$$

Proof. Since $\mathcal{H}_n(x) = \mathcal{P}_n(x-1)$, by Theorem 3.4, we easily obtain that if n is even, then

$$\mathcal{H}_{n+1}(x) = x\mathcal{H}_n(x),$$

and if n is odd, then

$$(x-1)\mathcal{H}_{n+1}(x) = x\mathcal{H}_n(x) - \frac{2}{n+1} \binom{n-1}{\frac{n-1}{2}},$$

for any $n \geq 3$, with initial condition $\mathcal{H}_3(x) = x$.

Since $\mathcal{H}(x, y) = \mathcal{P}(x-1, y)$, we have

$$\mathcal{H}(x, y) = \frac{[(x^2 - 1)y^2 - (x-1)y^2 C(y^2)](1+xy)}{x(x-1-xy^2)} - y^2.$$

■

Corollary 3.8 *Let the sequence $(h_{n,0}, h_{n,1}, \dots, h_{n, \lfloor \frac{1}{2}(n-1) \rfloor})$ be h -vector of $\mathcal{P}_n$. Then the sequence $h_{n,i}$ satisfies the following recurrence relation:*

$$\begin{cases} h_{n+1,0} &= h_{n,0} \\ h_{n+1,i} &= h_{n,i} + \varepsilon(n)h_{n+1,i-1} \quad \text{if } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1 \\ h_{n+1, \lfloor \frac{n}{2} \rfloor} &= \varepsilon(n)c_{\lfloor \frac{n}{2} \rfloor} \end{cases}$$

where $c_m = \frac{1}{m+1} \binom{2m}{m}$ and $\varepsilon(n) = 0$ if n is even; otherwise, $\varepsilon(n) = 1$, with initial conditions $(h_{3,0}, h_{3,1}) = (1, 0)$. Equivalently,

$$h_{n,i} = \frac{\lfloor \frac{n}{2} \rfloor - i}{\lfloor \frac{n}{2} \rfloor + i} \binom{\lfloor \frac{n}{2} \rfloor + i}{\lfloor \frac{n}{2} \rfloor}.$$

Proof. By comparing with the coefficient in Corollary 3.7, we can obtain the desired recurrence relations. Consider $t_{n,i} = \frac{\lfloor \frac{n}{2} \rfloor - i}{\lfloor \frac{n}{2} \rfloor + i} \binom{\lfloor \frac{n}{2} \rfloor + i}{\lfloor \frac{n}{2} \rfloor}$. Note that $t_{n,i}$ satisfies the above recurrence relations as well. Hence,

$$h_{n,i} = t_{n,i} = \frac{\lfloor \frac{n}{2} \rfloor - i}{\lfloor \frac{n}{2} \rfloor + i} \binom{\lfloor \frac{n}{2} \rfloor + i}{\lfloor \frac{n}{2} \rfloor}.$$

■

Remark 3.1 *Let $n \geq 3$. The number of left factor of Dyck path from $(0, 0)$ to $(\lfloor \frac{n}{2} \rfloor + i - 1, \lfloor \frac{n}{2} \rfloor - i - 1)$ equals $\frac{\lfloor \frac{n}{2} \rfloor - i}{\lfloor \frac{n}{2} \rfloor + i} \binom{\lfloor \frac{n}{2} \rfloor + i}{\lfloor \frac{n}{2} \rfloor}$ for any $0 \leq i \leq \lfloor \frac{n-1}{2} \rfloor$.*

4 The Algebras $\mathcal{A}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n}$ from the Poset $\mathcal{P}_n$

In this section, we will consider the properties of the algebras $\mathcal{A}_{\mathcal{P}_n}$ and $\mathcal{B}_{\mathcal{P}_n}$.

Let $m = \binom{n-1}{\lfloor \frac{1}{2}(n-1) \rfloor}$. We list all the elements of $\mathcal{P}_n$ as $S_{n,1}, S_{n,2}, \dots, S_{n,m}$. For any a sequence $S_{n,j_1}, S_{n,j_2}, \dots, S_{n,j_i}$ in the poset $\mathcal{P}_n$, let $\mathbf{r}(S_{n,j_1}, S_{n,j_2}, \dots, S_{n,j_i}) = (r_1, \dots, r_m)$ be a vector such that $r_j = |\{k \mid S_{n,j_k} = S_{n,j}\}|$. Furthermore, we can obtain a monomial

$$m(S_{n,j_1}, S_{n,j_2}, \dots, S_{n,j_i}) = x_1^{r_1} x_2^{r_2} \cdots x_m^{r_m}.$$

A sequence $S_{n,j_1}, S_{n,j_2}, \dots, S_{n,j_i}$ of elements in the poset $\mathcal{P}_n$ forms a multichain if and only if the monomial $m(S_{n,j_1}, S_{n,j_2}, \dots, S_{n,j_i})$ is nonvanishing in the algebra $\mathcal{A}_{\mathcal{P}_n}$.

For a monomial ideal $\mathcal{I}$, the set of all monomials that do not belong to $\mathcal{I}$ is a basis of the quotient of the polynomial ring modulo $\mathcal{I}$, called the standard monomial basis. Thus the monomials $m(S_{n,j_1}, S_{n,j_2}, \dots, S_{n,j_i})$, where $(S_{n,j_1}, S_{n,j_2}, \dots, S_{n,j_i})$ ranges over the multichains of the poset $\mathcal{P}_n$, form the standard monomial basis of the algebra $\mathcal{A}_{\mathcal{P}_n}$. The algebra $\mathcal{A}_{\mathcal{P}_n}$ is graded. For a graded algebra $\mathcal{A} = \mathcal{A}^0 \oplus \mathcal{A}^1 \oplus \mathcal{A}^2 \dots$, the Hilbert series of the algebra $\mathcal{A}$, is the formal power series in x given by

$$\text{Hilb}\mathcal{A} = \sum_{i \geq 0} x^i \dim \mathcal{A}^i;$$

there exists a polynomial $P_{\mathcal{A}}(x)$ with rational coefficients (called the *Hilbert polynomial* of $\mathcal{A}$) such that $P_{\mathcal{A}}(k) = \dim \mathcal{A}^k$ for all sufficiently large k .

Theorem 4.1 *The Hilbert polynomial $P_{\mathcal{A}_{\mathcal{P}_n}}(x)$ of the algebra $\mathcal{A}_{\mathcal{P}_n}$ is $x^{\lfloor \frac{1}{2}(n-1) \rfloor} \mathcal{P}_n(\frac{1}{x})$.*

Proof. Note that the number of the multichains $S_{n,j_1} \preceq S_{n,j_2} \preceq \dots \preceq S_{n,j_i}$ in $\mathcal{P}_n$ is equal to the dimension of $\mathcal{A}_n^i$. Corollary 3.5 implies that $\dim \mathcal{A}_n^i = i^{\lfloor \frac{n-1}{2} \rfloor} \mathcal{P}_n(\frac{1}{i})$ for any $i \geq 1$. Since $\deg(\mathcal{P}_n(x)) = \lfloor \frac{n-1}{2} \rfloor$, we have $P_{\mathcal{A}_{\mathcal{P}_n}}(x) = x^{\lfloor \frac{1}{2}(n-1) \rfloor} \mathcal{P}_n(\frac{1}{x})$. ■

Theorem 4.2 *For any $n \geq 3$, the Hilbert series $\text{Hilb}\mathcal{A}_{\mathcal{P}_n}(x)$ of the algebra $\mathcal{A}_{\mathcal{P}_n}$ satisfies the following recurrence: when n is even,*

$$\text{Hilb}\mathcal{A}_{\mathcal{P}_{n+1}}(x) = x\text{Hilb}\mathcal{A}'_{\mathcal{P}_n}(x) + \text{Hilb}\mathcal{A}_{\mathcal{P}_n}(x);$$

when n is odd,

$$\begin{aligned} \text{Hilb}\mathcal{A}_{\mathcal{P}_{n+3}}(x) &= x\text{Hilb}\mathcal{A}'_{\mathcal{P}_{n+2}}(x) + \text{Hilb}\mathcal{A}_{\mathcal{P}_{n+2}}(x) + \frac{4n}{n+3}x\text{Hilb}\mathcal{A}'_{\mathcal{P}_{n+1}}(x) \\ &\quad - \frac{8n}{n+3}x\text{Hilb}\mathcal{A}'_{\mathcal{P}_n}(x) - \frac{4n}{n+3}x^2\text{Hilb}\mathcal{A}''_{\mathcal{P}_n}(x), \end{aligned}$$

where the notation “ ’ ” denotes differentiation of functions, with the initial condition $\text{Hilb}\mathcal{A}_{\mathcal{P}_3}(x) = \frac{1}{(1-x)^2}$ and $\text{Hilb}\mathcal{A}_{\mathcal{P}_4}(x) = \frac{1+x}{(1-x)^2}$.

Proof. By Theorem 4.1, we have $\text{Hilb}\mathcal{A}_{\mathcal{P}_n}(x) = 1 + \sum_{i \geq 1} i^{\lfloor \frac{n-1}{2} \rfloor} \mathcal{P}_n(\frac{1}{i})x^i$. The results follow Theorem 3.4. ■

In general, we may suppose that $\text{Hilb}\mathcal{A}_{\mathcal{P}_n}(x) = \frac{A_{\mathcal{P}_n}(x)}{(1-x)^{\lfloor \frac{n-1}{2} \rfloor}}$, where $A_{\mathcal{P}_n}(x)$ is a polynomial.

Corollary 4.1 *For any $n \geq 3$, $A_{\mathcal{P}_n}(x)$ satisfies the following recurrence: when n is even,*

$$A_{\mathcal{P}_{n+1}}(x) = x(1-x)A'_{\mathcal{P}_n}(x) + \left[\left(\frac{1}{2}n - 1 \right)x + 1 \right] A_{\mathcal{P}_n}(x);$$

when n is odd,

$$\begin{aligned} (1-x)A_{\mathcal{P}_{n+3}}(x) &= x(1-x)A'_{\mathcal{P}_{n+2}}(x) + \frac{(n+1)x+2}{2}A_{\mathcal{P}_{n+2}}(x) + \frac{4n}{n+3}x(1-x)^2A'_{\mathcal{P}_{n+1}}(x) \\ &\quad + \frac{2n(n+1)}{n+3}x(1-x)A_{\mathcal{P}_{n+1}}(x) - \frac{4n}{n+3}x(1-x)[(n-1)x+2]A'_{\mathcal{P}_n}(x) \\ &\quad - \frac{n(n+1)}{n+3}x[(n-1)x+4]A_{\mathcal{P}_n}(x) - \frac{4n}{n+3}x^2(1-x)^2A''_{\mathcal{P}_n}(x) \end{aligned}$$

with the initial conditions $A_{\mathcal{P}_3}(x) = 1$ and $A_{\mathcal{P}_4}(x) = 1 + x$.

Proof. By Theorem 4.2, we immediately obtain the desired results after simple computations. ■

Note that a sequence $S_{n,i_1}, S_{n,i_2}, \dots, S_{n,i_j}$ of elements in the poset $\mathcal{P}_n$ forms a chain if and only if the monomial $x_{i_1}x_{i_2} \cdots x_{i_j}$ is nonvanishing in the algebra $\mathcal{B}_{\mathcal{P}_n}$.

Theorem 4.3 *For any $n \geq 3$, the Hilbert series $\text{Hilb}\mathcal{B}_{\mathcal{P}_n}(x)$ of the algebra $\mathcal{B}_{\mathcal{P}_n}$ is*

$$\text{Hilb}\mathcal{B}_{\mathcal{P}_n}(x) = 1 + \sum_{i=1}^{\lfloor \frac{1}{2}(n-1) \rfloor + 1} \sum \binom{n}{d_1, d_2, \dots, d_{i+1}} \frac{2d_{i+1} - n}{n} x^i,$$

where the second sum is over all $(d_1, \dots, d_{i+1})$ such that $\sum_{k=1}^{i+1} d_k = n$, $d_1 \geq 0$, $d_k \geq 1$ for all $2 \leq k \leq i$ and $d_{i+1} \geq n - \lfloor \frac{n-1}{2} \rfloor$.

Proof. Note that the number of the chains $S_{n,j_1} \prec S_{n,j_2} \prec \cdots \prec S_{n,j_i}$ in $\mathcal{P}_n$ is equal to the dimension of $\mathcal{B}_{\mathcal{P}_n}^i$. Theorem 3.5 implies that $\dim \mathcal{B}_{\mathcal{P}_n}^i = \sum \binom{n}{d_1, d_2, \dots, d_{i+1}} \frac{2d_{i+1} - n}{n}$ for any

$i \geq 1$, where the sum is over all $(d_1, \dots, d_{k+1})$ such that $\sum_{k=1}^{i+1} d_k = n$, $d_1 \geq 0$, $d_k \geq 1$ for all $2 \leq k \leq i$ and $d_{i+1} \geq n - \lfloor \frac{n-1}{2} \rfloor$. This completes the proof. ■

References

- [1] Hungyung Chang, Jun Ma, Yeong-Nan Yeh, Enumerations of Permutations by Circular Descent Sets
- [2] Robert J. Clarke, Einar Steingrímsson, Jiang Zeng, New Euler-Mahonian permutation statistics, *Sm. Lothar. Combin. 35 (1995), Art. B35c, approx. 29 pp. (electronic)*.
- [3] Michael Domaratzki, Combinatorial Interpretations of a Generalization of the Genocchi Numbers, *Journal of Integer Sequences, Vol. 7 (2004), Article 04.3.6*
- [4] Cori, R. and X. G. Viennot, A synthesis of bijections related to Catalan numbers, unpublished, 1983.
- [5] Richard P. Stanley, Enumerative Combinatorics, Cambridge ; New York : Cambridge University Press, c1997-1999
- [6] J. R. Stembridge, Enriched P-partitions, *Trans. Amer. Math. Soc.* 349 (1997), no. 2, 763-788.